

CLASS XII

CHAPTER 5

Theorem 5 (To be inserted on page 173 under the heading theorem 5)

(i) Derivative of Exponential Function $f(x) = e^x$.

If $f(x) = e^x$, then

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{e^{x+\Delta x} - e^x}{\Delta x} \\ &= e^x \cdot \lim_{\Delta x \rightarrow 0} \frac{e^{\Delta x} - 1}{\Delta x} \\ &= e^x \cdot 1 \text{ [since } \lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1 \text{]} \end{aligned}$$

Thus, $\frac{d}{dx}(e^x) = e^x$.

(ii) Derivative of logarithmic function $f(x) = \log_e x$.

If $f(x) = \log_e x$, then

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{\log_e(x + \Delta x) - \log_e x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{\log_e \left(1 + \frac{\Delta x}{x}\right)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{x} \frac{\log_e \left(1 + \frac{\Delta x}{x}\right)}{\frac{\Delta x}{x}} \\ &= \frac{1}{x} \text{ [since } \lim_{h \rightarrow 0} \frac{\log_e(1+h)}{h} = 1 \text{]} \end{aligned}$$

Thus, $\frac{d}{dx} \log_e x = \frac{1}{x}$

CHAPTER 7

7.6.3. $\int (px + q)\sqrt{ax^2 + bx + c} \, dx.$

We choose constants A and B such that

$$\begin{aligned} px + q &= A \left[\frac{d}{dx}(ax^2 + bx + c) \right] + B \\ &= A(2ax + b) + B \end{aligned}$$

Comparing the coefficients of x and the constant terms on both sides, we get

$$2aA = p \text{ and } Ab + B = q$$

Solving these equations, the values of A and B are obtained. Thus, the integral reduces to

$$\begin{aligned} A \int (2ax + b)\sqrt{ax^2 + bx + c} \, dx + B \int \sqrt{ax^2 + bx + c} \, dx \\ = AI_1 + BI_2 \end{aligned}$$

where

$$I_1 = \int (2ax + b)\sqrt{ax^2 + bx + c} \, dx$$

Put $ax^2 + bx + c = t$, then $(2ax + b)dx = dt$

So
$$I_1 = \frac{2}{3}(ax^2 + bx + c)^{\frac{3}{2}} + C_1$$

Similarly,

$$I_2 = \int \sqrt{ax^2 + bx + c} \, dx$$

is found, using the integral formula discussed in [7.6.2, Page 328 of the textbook].

Thus $\int (px + q)\sqrt{ax^2 + bx + c} \, dx$ is finally worked out.

Example 25 Find $\int x\sqrt{1+x-x^2} \, dx$

Solution Following the procedure as indicated above, we write

$$\begin{aligned} x &= A \left[\frac{d}{dx}(1 + x - x^2) \right] + B \\ &= A(1 - 2x) + B \end{aligned}$$

Equating the coefficients of x and constant terms on both sides,

We get $-2A = 1$ and $A + B = 0$

Solving these equations, we get $A = -\frac{1}{2}$ and $B = \frac{1}{2}$. Thus the integral reduces to

$$\begin{aligned}\int x\sqrt{1+x-x^2} dx &= -\frac{1}{2}\int(1-2x)\sqrt{1+x-x^2} dx + \frac{1}{2}\int\sqrt{1+x-x^2} dx \\ &= -\frac{1}{2}I_1 + \frac{1}{2}I_2\end{aligned}\quad (1)$$

Consider $I_1 = \int(1-2x)\sqrt{1+x-x^2} dx$

Put $1+x-x^2 = t$, then $(1-2x)dx = dt$

$$\begin{aligned}\text{Thus } I_1 &= \int(1-2x)\sqrt{1+x-x^2} dx = \int t^{\frac{1}{2}} dt = \frac{2}{3}t^{\frac{3}{2}} + C_1 \\ &= \frac{2}{3}(1+x-x^2)^{\frac{3}{2}} + C_1, \text{ where } C_1 \text{ is some constant.}\end{aligned}$$

Further, consider $I_2 = \int\sqrt{1+x-x^2} dx = \int\sqrt{\frac{5}{4} - \left(x - \frac{1}{2}\right)^2} dx$

Put $x - \frac{1}{2} = t$. Then $dx = dt$

$$\begin{aligned}\text{Therefore, } I_2 &= \int\sqrt{\left(\frac{\sqrt{5}}{2}\right)^2 - t^2} dt \\ &= \frac{1}{2}t\sqrt{\frac{5}{4} - t^2} + \frac{1}{2} \cdot \frac{5}{4} \sin^{-1} \frac{2t}{\sqrt{5}} + C_2 \\ &= \frac{1}{2} \frac{(2x-1)}{2} \sqrt{\frac{5}{4} - \left(x - \frac{1}{2}\right)^2} + \frac{5}{8} \sin^{-1} \left(\frac{2x-1}{\sqrt{5}}\right) + C_2 \\ &= \frac{1}{4}(2x-1)\sqrt{1+x-x^2} + \frac{5}{8} \sin^{-1} \left(\frac{2x-1}{\sqrt{5}}\right) + C_2, \text{ where } C_2\end{aligned}$$

is some constant.

Putting values of I_1 and I_2 in (1), we get

$$\begin{aligned}\int x\sqrt{1+x-x^2} dx &= -\frac{1}{3}(1+x-x^2)^{\frac{3}{2}} + \frac{1}{8}(2x-1)\sqrt{1+x-x^2} \\ &\quad + \frac{5}{16} \sin^{-1} \left(\frac{2x-1}{\sqrt{5}}\right) + C,\end{aligned}$$

where

$$C = -\frac{C_1 + C_2}{2} \text{ is another arbitrary constant.}$$

Insert the following exercises at the end of EXERCISE 7.7 as follows:

12. $x\sqrt{x+x^2}$ 13. $(x+1)\sqrt{2x^2+3}$ 14. $(x+3)\sqrt{3-4x-x^2}$

Answers

12. $\frac{1}{3}(x^2+x)^{\frac{3}{2}} - \frac{(2x+1)\sqrt{x^2+x}}{8} + \frac{1}{16} \log |x + \frac{1}{2} + \sqrt{x^2+x}| + C$

13. $\frac{1}{6}(2x^2+3)^{\frac{3}{2}} + \frac{x}{2}\sqrt{2x^2+3} + \frac{3\sqrt{2}}{4} \log \left| x + \sqrt{x^2 + \frac{3}{2}} \right| + C$

14. $-\frac{1}{3}(3-4x-x^2)^{\frac{3}{2}} + \frac{7}{2} \sin^{-1} \left(\frac{x+2}{\sqrt{7}} \right) + \frac{(x+2)\sqrt{3-4x-x^2}}{2} + C$

CHAPTER 10

10.7 Scalar triple product Let \vec{a}, \vec{b} and \vec{c} be any three vectors. The scalar product of \vec{a} and $(\vec{b} \times \vec{c})$, i.e., $\vec{a} \cdot (\vec{b} \times \vec{c})$ is called the scalar triple product of \vec{a}, \vec{b} and \vec{c} in this order and is denoted by $[\vec{a}, \vec{b}, \vec{c}]$ (or $[\vec{a} \vec{b} \vec{c}]$). We thus have

$$[\vec{a}, \vec{b}, \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c})$$

Observations

1. Since $(\vec{b} \times \vec{c})$ is a vector, $\vec{a} \cdot (\vec{b} \times \vec{c})$ is a scalar quantity, i.e. $[\vec{a}, \vec{b}, \vec{c}]$ is a scalar quantity.

2. Geometrically, the magnitude of the scalar triple product is the volume of a parallelepiped formed by adjacent sides given by the three vectors \vec{a}, \vec{b} and \vec{c} (Fig. 10.28). Indeed, the area of the parallelogram forming the base of the parallelepiped is $|\vec{b} \times \vec{c}|$. The height is the projection of \vec{a} along the normal to the plane containing \vec{b} and \vec{c} which is the magnitude of the component of \vec{a} in the direction of $\vec{b} \times \vec{c}$




Fig. 10.28

i.e., $\frac{|\vec{a} \cdot (\vec{b} \times \vec{c})|}{|\vec{b} \times \vec{c}|}$. So the required volume of the parallelepiped

is $\frac{|\vec{a} \cdot (\vec{b} \times \vec{c})|}{|\vec{b} \times \vec{c}|} |\vec{b} \times \vec{c}| = |\vec{a} \cdot (\vec{b} \times \vec{c})|$,

3. If $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$, $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ and $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$, then

$$\hat{b} \times \hat{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$= (b_2c_3 - b_3c_2) \hat{i} + (b_3c_1 - b_1c_3) \hat{j} + (b_1c_2 - b_2c_1) \hat{k}$$

and so

$$\hat{a} \cdot (\hat{b} \times \hat{c}) = a_1(b_2c_3 - b_3c_2) + a_2(b_3c_1 - b_1c_3) + a_3(b_1c_2 - b_2c_1)$$

$$= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

4. If \hat{a}, \hat{b} and \hat{c} be any three vectors, then

$$[\hat{a}, \hat{b}, \hat{c}] = [\hat{b}, \hat{c}, \hat{a}] = [\hat{c}, \hat{a}, \hat{b}]$$

(cyclic permutation of three vectors does not change the value of the scalar triple product).

Let $\hat{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$, $\hat{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ and $\hat{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$. Then, just by observation above, we have

$$\begin{aligned} [\hat{a}, \hat{b}, \hat{c}] &= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= a_1(b_2c_3 - b_3c_2) + a_2(b_3c_1 - b_1c_3) + a_3(b_1c_2 - b_2c_1) \\ &= b_1(a_3c_2 - a_2c_3) + b_2(a_1c_3 - a_3c_1) + b_3(a_2c_1 - a_1c_2) \\ &= \begin{vmatrix} b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \\ a_1 & a_2 & a_3 \end{vmatrix} \\ &= [\hat{b}, \hat{c}, \hat{a}] \end{aligned}$$

Similarly, the reader may verify that

$$= [\hat{a}, \hat{b}, \hat{c}] = [\hat{c}, \hat{a}, \hat{b}]$$

Hence $[\hat{a}, \hat{b}, \hat{c}] = [\hat{b}, \hat{c}, \hat{a}] = [\hat{c}, \hat{a}, \hat{b}]$

5. In scalar triple product $\hat{a} \cdot (\hat{b} \times \hat{c})$, the dot and cross can be interchanged. Indeed,

$$\hat{a} \cdot (\hat{b} \times \hat{c}) = [\hat{a}, \hat{b}, \hat{c}] = [\hat{b}, \hat{c}, \hat{a}] = [\hat{c}, \hat{a}, \hat{b}] = \hat{c} \cdot (\hat{a} \times \hat{b}) = (\hat{a} \times \hat{b}) \cdot \hat{c}$$

6. $[\hat{a}, \hat{b}, \hat{c}] = -[\hat{a}, \hat{c}, \hat{b}]$. Indeed

$$\begin{aligned}
 &= [\mathbf{a}, \mathbf{b}, \mathbf{c}] = \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) \\
 &= \mathbf{a} \cdot (-\mathbf{c} \times \mathbf{b}) \\
 &= -(\mathbf{a} \cdot (\mathbf{c} \times \mathbf{b})) \\
 &= -[\mathbf{a}, \mathbf{c}, \mathbf{b}]
 \end{aligned}$$

7. $[\mathbf{a}, \mathbf{a}, \mathbf{b}] = 0$. Indeed

$$\begin{aligned}
 [\mathbf{a}, \mathbf{a}, \mathbf{b}] &= [\mathbf{a}, \mathbf{b}, \mathbf{a}] \\
 &= [\mathbf{b}, \mathbf{a}, \mathbf{a}] \\
 &= \mathbf{b} \cdot (\mathbf{a} \times \mathbf{a}) \\
 &= \mathbf{b} \cdot \mathbf{0} = 0. \qquad \qquad \qquad (\text{as } \mathbf{a} \times \mathbf{a} = \mathbf{0})
 \end{aligned}$$

Note: The result in 7 above is true irrespective of the position of two equal vectors.

10.7.1 Coplanarity of three vectors

Theorem 1 Three vectors \mathbf{a} , \mathbf{b} and \mathbf{c} are coplanar if and only if $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = 0$.

Proof : Suppose first that the vectors \mathbf{a} , \mathbf{b} and \mathbf{c} are coplanar.

If \mathbf{b} and \mathbf{c} are parallel vectors, then, $\mathbf{b} \times \mathbf{c} = \mathbf{0}$ and so $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = 0$.

If \mathbf{b} and \mathbf{c} are not parallel then, since \mathbf{a} , \mathbf{b} and \mathbf{c} are coplanar, $\mathbf{b} \times \mathbf{c}$ is perpendicular to \mathbf{a} .

So $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = 0$.

Conversely, suppose that $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = 0$. If \mathbf{a} and $\mathbf{b} \times \mathbf{c}$ are both non-zero, then we conclude that \mathbf{a} and $\mathbf{b} \times \mathbf{c}$ are perpendicular vectors. But $\mathbf{b} \times \mathbf{c}$ is perpendicular to both \mathbf{b} and \mathbf{c} . Therefore \mathbf{a} and \mathbf{b} and \mathbf{c} must lie in the plane, i.e. they are coplanar. If $\mathbf{a} = \mathbf{0}$, then \mathbf{a} is coplanar with any two vectors, in particular with \mathbf{b} and \mathbf{c} . If $(\mathbf{b} \times \mathbf{c}) = \mathbf{0}$, then \mathbf{b} and \mathbf{c} are parallel vectors and so, \mathbf{a} , \mathbf{b} and \mathbf{c} are coplanar since any two vectors always lie in a plane determined by them and a vector which is parallel to any one of it also lies in that plane.

Note: Coplanarity of four points can be discussed using coplanarity of three vectors. Indeed, the four points A, B, C and D are coplanar if the vectors \overrightarrow{AB} , \overrightarrow{AC} and \overrightarrow{AD} are coplanar.

Example 26: Find $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$, if $\mathbf{a} = 2\hat{i} + \hat{j} + 3\hat{k}$, $\mathbf{b} = -\hat{i} + 2\hat{j} + \hat{k}$ and $\mathbf{c} = 3\hat{i} + \hat{j} + 2\hat{k}$.

Solution : We have $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 2 & 1 & 3 \\ -1 & 2 & 1 \\ 3 & 1 & 2 \end{vmatrix} = -10$.

Example 27: Show that the vectors $\mathbf{a} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\mathbf{b} = -2\hat{i} + 3\hat{j} - 4\hat{k}$ and $\mathbf{c} = \hat{i} - 3\hat{j} + 5\hat{k}$ are coplanar.

Solution : We have $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} 1 & -2 & 3 \\ -2 & 2 & -4 \\ 1 & -3 & 5 \end{vmatrix} = 0$.

Hence, in view of Theorem 1, \hat{a}, \hat{b} and \hat{c} are coplanar vectors.

Example 28: Find λ if the vectors $\hat{r} = \hat{i} + 3\hat{j} + \hat{k}$, $\hat{b} = 2\hat{i} - \hat{j} - \hat{k}$ and $\hat{c} = \lambda\hat{i} + 7\hat{j} + 3\hat{k}$ are coplanar.

Solution : Since \hat{a}, \hat{b} and \hat{c} are coplanar vectors, we have $[\hat{a}, \hat{b}, \hat{c}] = 0$, i.e.,

$$\begin{vmatrix} 1 & 3 & 1 \\ 2 & -1 & -1 \\ \lambda & 7 & 3 \end{vmatrix} = 0.$$

$$\Rightarrow 1(-3+7) - 3(6+\lambda) + 1(14+\lambda) = 0$$

$$\Rightarrow \lambda = 0.$$

Example 29: Show that the four points A, B, C and D with position vectors $4\hat{i} + 5\hat{j} + \hat{k}$, $-(\hat{j} + \hat{k})$, $3\hat{i} + 9\hat{j} + 4\hat{k}$ and $4(-\hat{i} + \hat{j} + \hat{k})$, respectively are coplanar.

Solution : We know that the four points A, B, C and D are coplanar if the three vectors $\overrightarrow{AB}, \overrightarrow{AC}$ and \overrightarrow{AD} are coplanar, i.e., if

$$[\overrightarrow{AB}, \overrightarrow{AC}, \overrightarrow{AD}] = 0$$

Now $\overrightarrow{AB} = -(\hat{j} + \hat{k}) - (4\hat{i} + 5\hat{j} + \hat{k}) = -4\hat{i} - 6\hat{j} - 2\hat{k}$

$$\overrightarrow{AC} = (3\hat{i} + 9\hat{j} + 4\hat{k}) - (4\hat{i} + 5\hat{j} + \hat{k}) = -\hat{i} + 4\hat{j} + 3\hat{k}$$

and $\overrightarrow{AD} = 4(-\hat{i} + \hat{j} + \hat{k}) - (4\hat{i} + 5\hat{j} + \hat{k}) = -8\hat{i} - \hat{j} + 3\hat{k}$

Thus $[\overrightarrow{AB}, \overrightarrow{AC}, \overrightarrow{AD}] = \begin{vmatrix} -4 & -6 & -2 \\ -1 & 4 & 3 \\ -8 & -1 & 3 \end{vmatrix} = 0.$

Hence A, B, C and D are coplanar.

Example 30 : Prove that $[\hat{a} + \hat{b}, \hat{b} + \hat{c}, \hat{c} + \hat{a}] = 2[\hat{a}, \hat{b}, \hat{c}]$.

Solution : We have

$$\begin{aligned} [\hat{a} + \hat{b}, \hat{b} + \hat{c}, \hat{c} + \hat{a}] &= (\hat{a} + \hat{b}) \cdot ((\hat{b} + \hat{c}) \times (\hat{c} + \hat{a})) \\ &= (\hat{a} + \hat{b}) \cdot (\hat{b} \times \hat{c} + \hat{b} \times \hat{a} + \hat{c} \times \hat{c} + \hat{c} \times \hat{a}) \\ &= (\hat{a} + \hat{b}) \cdot (\hat{b} \times \hat{c} + \hat{b} \times \hat{a} + \hat{c} \times \hat{a}) \quad (\text{as } \hat{c} \times \hat{c} = \hat{0}) \\ &= \hat{a} \cdot (\hat{b} \times \hat{c}) + \hat{a} \cdot (\hat{b} \times \hat{a}) + \hat{a} \cdot (\hat{c} \times \hat{a}) + \hat{b} \cdot (\hat{b} \times \hat{c}) + \hat{b} \cdot (\hat{b} \times \hat{a}) + \hat{b} \cdot (\hat{c} \times \hat{a}) \\ &= [\hat{a}, \hat{b}, \hat{c}] + [\hat{a}, \hat{b}, \hat{a}] + [\hat{a}, \hat{c}, \hat{a}] + [\hat{b}, \hat{b}, \hat{c}] + [\hat{b}, \hat{b}, \hat{a}] + [\hat{b}, \hat{c}, \hat{a}] \end{aligned}$$

$$= 2[\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{c}] \quad (\text{Why?})$$

Example 31 : Prove that $[\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{c} + \mathbf{d}] = [\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{c}] + [\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{d}]$

Solution We have

$$\begin{aligned} [\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{c} + \mathbf{d}] &= \mathbf{r} \cdot (\mathbf{b} \times (\mathbf{c} + \mathbf{d})) \\ &= \mathbf{r} \cdot (\mathbf{b} \times \mathbf{c} + \mathbf{b} \times \mathbf{d}) \\ &= \mathbf{r} \cdot (\mathbf{b} \times \mathbf{c}) + \mathbf{r} \cdot (\mathbf{b} \times \mathbf{d}) \\ &= [\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{c}] + [\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{d}] \end{aligned}$$

EXERCISE 10.5

- Find $[\mathbf{r}, \mathbf{a}, \mathbf{b}, \mathbf{c}]$ if $\mathbf{a} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\mathbf{b} = 2\hat{i} - 3\hat{j} + \hat{k}$ and $\mathbf{c} = 3\hat{i} + \hat{j} - 2\hat{k}$
(Answer : 24)
- Show that the vectors $\mathbf{a} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\mathbf{b} = -2\hat{i} + 3\hat{j} - 4\hat{k}$ and $\mathbf{c} = \hat{i} - 3\hat{j} + 5\hat{k}$ are coplanar.
- Find λ if the vectors $\hat{i} - \hat{j} + \hat{k}$, $3\hat{i} + \hat{j} + 2\hat{k}$ and $\hat{i} + \lambda\hat{j} - 3\hat{k}$ are coplanar. (Answer : $\lambda = 15$)
- Let $\mathbf{a} = \hat{i} + \hat{j} + \hat{k}$, $\mathbf{b} = \hat{i}$ and $\mathbf{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$ Then
 - If $c_1 = 1$ and $c_2 = 2$, find c_3 which makes \mathbf{a} , \mathbf{b} and \mathbf{c} coplanar (Answer : $c_3 = 2$)
 - If $c_2 = -1$ and $c_3 = 1$, show that no value of c_1 can makes \mathbf{a} , \mathbf{b} and \mathbf{c} coplanar.
- Show that the four points with position vectors $4\hat{i} + 8\hat{j} + 12\hat{k}$, $2\hat{i} + 4\hat{j} + 6\hat{k}$, $3\hat{i} + 5\hat{j} + 4\hat{k}$ and $5\hat{i} + 8\hat{j} + 5\hat{k}$ are coplanar.
- Find x such that the four points A (3, 2, 1) B (4, x, 5), C (4, 2, -2) and D (6, 5, -1) are coplanar.
(Answer $x = 5$)
- Show that the vectors \mathbf{a} , \mathbf{b} and \mathbf{c} coplanar if $\mathbf{a} + \mathbf{b}$, $\mathbf{b} + \mathbf{c}$ and $\mathbf{c} + \mathbf{a}$ are coplanar.

MATHEMATICS



PioneerMathematics.com