

## JEE-Main Mathematics Questions Paper 18.3.2021 Shift-1


1. The differential equation satisfied by the system of parabolas  $y^2 = 4a(x + a)$  is:
- (a)  $y\left(\frac{dy}{dx}\right)^2 - 2x\left(\frac{dy}{dx}\right) - y = 0$
- (b)  $y\left(\frac{dy}{dx}\right)^2 - 2x\left(\frac{dy}{dx}\right) + y = 0$
- (c)  $y\left(\frac{dy}{dx}\right)^2 + 2x\left(\frac{dy}{dx}\right) - y = 0$
- (d)  $y\left(\frac{dy}{dx}\right)^2 + 2x\left(\frac{dy}{dx}\right) + y = 0$
2. The number of integral values of  $m$  so that the abscissa of point of intersection of lines  $3x + 4y = 9$  and  $y = mx + 1$  is also an integer, is:
- (a) 1 (b) 2 (c) 3 (d) 0
3. Let  $(1 + x + 2x^2)^{20} = a_0 + a_1x + a_2x^2 + \dots + a_{40}x^{40}$ . Then  $a_1 + a_3 + a_5 + \dots + a_{37}$  is equal to
- (a)  $2^{20}(2^{20} - 21)$  (b)  $2^{19}(2^{20} - 21)$  (c)  $2^{19}(2^{20} + 21)$  (d)  $2^{20}(2^{20} + 21)$
4. The solutions of the equation
- $$\begin{vmatrix} 1 + \sin^2 x & \sin^2 x & \sin^2 x \\ \cos^2 x & 1 + \cos^2 x & \cos^2 x \\ 4\sin 2x & 4\sin 2x & 1 + 4\sin 2x \end{vmatrix} = 0, (0 < x < \pi), \text{ are}$$
- (a)  $\frac{\pi}{12}, \frac{\pi}{6}$  (b)  $\frac{\pi}{6}, \frac{5\pi}{6}$  (c)  $\frac{5\pi}{12}, \frac{7\pi}{12}$  (d)  $\frac{7\pi}{12}, \frac{11\pi}{12}$
5. Choose the correct statement about two circles whose equations are given below:
- $$x^2 + y^2 - 10x - 10y + 41 = 0$$
- $$x^2 + y^2 - 22x - 10y + 137 = 0$$
- (a) circles have same centre (b) circles have no meeting point
- (c) circles have only one meeting point (d) circles have two meeting points
6. Let  $\alpha, \beta, \gamma$  be the real roots of the equation,  $x^3 + ax^2 + bx + c = 0$ , ( $a, b, c \in \mathbb{R}$  and  $a, b \neq 0$ ). If the system of equations (in  $u, v, w$ ) given by  $\alpha u + \beta v + \gamma w = 0$ ,  $\beta u + \gamma v + \alpha w = 0$ ;  $\gamma u + \alpha v + \beta w = 0$  has non-trivial solution, then the value of  $\frac{a^2}{b}$  is
- (a) 5 (b) 3 (c) 1 (d) 0
7. The integral  $\int \frac{(2x-1)\cos\sqrt{(2x-1)^2+5}}{\sqrt{4x^2-4x+6}} dx$  is equal to
- (where  $c$  is a constant of integration)
- (a)  $\frac{1}{2}\sin\sqrt{(2x-1)^2+5} + c$  (b)  $\frac{1}{2}\cos\sqrt{(2x+1)^2+5} + c$
- (c)  $\frac{1}{2}\cos\sqrt{(2x-1)^2+5} + c$  (d)  $\frac{1}{2}\sin\sqrt{(2x+1)^2+5} + c$

8. The equation of one of the straight lines which passes through the point (1, 3) and makes an angle  $\tan^{-1}(\sqrt{2})$  with the straight line,  $y + 1 = 3\sqrt{2}x$  is
- (a)  $4\sqrt{2}x + 5y - (15 + 4\sqrt{2}) = 0$       (b)  $5\sqrt{2}x + 4y - (15 + 4\sqrt{2}) = 0$   
(c)  $4\sqrt{2}x + 5y - 4\sqrt{2} = 0$       (d)  $4\sqrt{2}x - 5y - (5 + 4\sqrt{2}) = 0$
9. If  $\lim_{x \rightarrow 0} \frac{\sin^{-1} x - \tan^{-1} x}{3x^3}$  is equal to L, then the value of  $(6L + 1)$  is
- (a)  $\frac{1}{6}$       (b)  $\frac{1}{2}$       (c) 6      (d) 2
10. A vector  $\vec{a}$  has components  $3p$  and  $1$  with respect to a rectangular cartesian system. This system is rotated through a certain angle about the origin in the counter clockwise sense. If, with respect to new system,  $\vec{a}$  has components  $p + 1$  and  $\sqrt{10}$ , then a value of  $p$  is equal to:
- (a) 1      (b)  $-\frac{5}{4}$       (c)  $\frac{4}{5}$       (d) -1
11. If the equation  $a|z|^2 + \overline{\alpha z} + \alpha \overline{z} + d = 0$  represents a circle where  $a, d$  are real constants then which of the following condition is correct?
- (a)  $|\alpha|^2 - ad \neq 0$   
(b)  $|\alpha|^2 - ad > 0$  and  $a \in \mathbb{R} - \{0\}$   
(c)  $|\alpha|^2 - ad \geq 0$  and  $a \in \mathbb{R}$   
(d)  $\alpha = 0, a, d \in \mathbb{R}^+$
12. For the four circles M, N, O and P, following four equations are given:  
Circle M :  $x^2 + y^2 = 1$   
Circle N :  $x^2 + y^2 - 2x = 0$   
Circle O :  $x^2 + y^2 - 2x - 2y + 1 = 0$   
Circle P :  $x^2 + y^2 - 2y = 0$   
If the centre of circle M is joined with centre of the circle N, further centre of circle N is joined with centre of the circle O, centre of circle O is joined with the centre of circle P and lastly, centre of circle P is joined with centre of circle M, then these lines from the sides of a:
- (a) Rhombus  
(b) Square  
(c) Rectangle  
(d) Parallelogram
13. If  $\alpha, \beta$  are natural numbers such that  $100^\alpha - 199\beta = (100)(100) + (99)(101) + (98)(102) + \dots + (1)(199)$ , then the slope of the line passing through  $(\alpha, \beta)$  and origin is:
- (a) 540      (b) 550      (c) 530      (d) 510

14. The real valued function  $f(x) = \frac{\operatorname{cosec}^{-1}x}{\sqrt{x-[x]}}$ , where  $[x]$  denotes the greatest integer less than or equal to  $x$ , is defined for all  $x$  belonging to:
- (a) all reals except integers  
 (b) all non-integers except the interval  $[-1, 1]$   
 (c) all integers except  $0, -1, 1$   
 (d) all reals except the interval  $[-1, 1]$
15.  $\frac{1}{3^2-1} + \frac{1}{5^2-1} + \frac{1}{7^2-1} + \dots + \frac{1}{(201)^2-1}$  is equal to
- (a)  $\frac{101}{404}$                       (b)  $\frac{25}{101}$                       (c)  $\frac{101}{408}$                       (d)  $\frac{99}{400}$
16. If the functions are defined as  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{1-x}$ , then what is the common domain of the following functions:  
 $f+g, f-g, f/g, g/f, g-f$  where  $(f \pm g)(x) = f(x) \pm g(x), (f/g)(x) = \frac{f(x)}{g(x)}$
- (a)  $0 \leq x \leq 1$                       (b)  $0 \leq x < 1$                       (c)  $0 < x < 1$                       (d)  $0 < x \leq 1$
17. If  $f(x) = \begin{cases} \frac{1}{|x|} & ; |x| \geq 1 \\ ax^2 + b & ; |x| < 1 \end{cases}$  is differentiable at every point of the domain, then the values of  $a$  and  $b$  are respectively:
- (a)  $\frac{1}{2}, \frac{1}{2}$                       (b)  $\frac{1}{2}, -\frac{3}{2}$                       (c)  $\frac{5}{2}, -\frac{3}{2}$                       (d)  $-\frac{1}{2}, \frac{3}{2}$
18. Let  $A + 2B = \begin{bmatrix} 1 & 2 & 0 \\ 6 & -3 & 3 \\ -5 & 3 & 1 \end{bmatrix}$  and  $2A - B = \begin{bmatrix} 2 & -1 & 5 \\ 2 & -1 & 6 \\ 0 & 1 & 2 \end{bmatrix}$ . If  $\operatorname{Tr}(A)$  denotes the sum of all diagonal elements of the matrix  $A$ , then  $\operatorname{Tr}(A) - \operatorname{Tr}(B)$  has value equal to
- (a) 1                      (b) 2                      (c) 0                      (d) 3
19. The sum of all the 4-digit distinct numbers that can be formed with the digits 1, 2, 2 and 3 is:
- (a) 26664                      (b) 122664                      (c) 122234                      (d) 22264
20. The value of  $3 + \frac{1}{4 + \frac{1}{3 + \frac{1}{4 + \frac{1}{3 + \dots \infty}}}}$  is equal to
- (a)  $1.5 + \sqrt{3}$                       (b)  $2 + \sqrt{3}$                       (c)  $3 + 2\sqrt{3}$                       (d)  $4 + \sqrt{3}$

**Integer Type**

- The number of times the digit 3 will be written when listing the integers from 1 to 1000 is
- Let the plane  $ax + by + cz + d = 0$  bisect the line joining the points  $(4, -3, 1)$  and  $(2, 3, -5)$  at the right angles. If  $a, b, c$  are integers, then the minimum value of  $(a^2 + b^2 + c^2 + d^2)$  is
- Let  $f(x)$  and  $g(x)$  be two functions satisfying  $f(x^2) + g(4 - x) = 4x^3$  and  $g(4 - x) + g(x) = 0$ , then the value of  $\int_{-4}^4 f(x)^2 dx$  is
- The missing value in the following figure is



- Let  $z_1, z_2$  be the roots of the equation  $z^2 + az + 12 = 0$  and  $z_1, z_2$  from an equilateral triangle with origin. Then, the value of  $|a|$  is
- The equation of the planes parallel to the plane  $x - 2y + 2z - 3 = 0$  which are at unit distance from the point  $(1, 2, 3)$  is  $ax + by + cz + d = 0$ . If  $(b - d) = K(c - a)$ , then the positive value of  $K$  is
- The mean age of 25 teachers in a school is 40 years. A teacher retires at the age of 60 years and a new teacher is appointed in his place. If the mean age of the teachers in this school now is 39 years, then the age (in years) of the newly appointed teacher is \_\_\_\_.
- If  $f(x) = \int \frac{5x^8 + 7x^6}{(x^2 + 1 + 2x^7)^2} dx, (x \geq 0), f(0) = 0$  and  $f(1) = \frac{1}{K}$ , then the value of  $K$  is
- A square ABCD has all its vertices on the curve  $x^2y^2 = 1$ . The mid-points of its sides also lie on the same curve. Then, the square of area of ABCD is
- The number of solutions of the equation  $|\cot x| = \cot x + \frac{1}{\sin x}$  in the interval  $[0, 2\pi]$  is

**ANSWER KEYS**

**Single Correct Type**

1.	c	2.	b	3.	b	4.	d	5.	c
6.	b	7.	a	8.	a	9.	d	10.	d
11.	b	12.	b	13.	b	14.	b	15.	b
16.	c	17.	d	18.	b	19.	a	20.	a

**Integer Type**

- |        |       |        |       |       |
|--------|-------|--------|-------|-------|
| 1. 300 | 2. 28 | 3. 512 | 4. 4  | 5. 35 |
| 6. 4   | 7. 35 | 8. 4   | 9. 80 | 10. 1 |